

Trigonometrical sums connected with one-dimensional lattice, the chiral Potts model, and number theory using the residue operator

NOUREDDINE CHAIR

Physics Department, University of Jordan, Amman, Jordan

Email: n.chair@ju.edu.jo

Abstract

We have recently developed a method for obtaining exact two-point resistance on the square N -cycle, the complete graph minus N edges of the opposite vertices, here, a similar method is used to obtain closed formulas of certain trigonometrical sums that arise in connection with one-dimensional lattice and the chiral Potts model. By modifying slightly, the previous computations, then, we are able to get new closed formulas for different trigonometrical sums, some of which appear in number theory.

1 Introduction

We developed Recently a method for computing exact two-point resistance of the square of the N -cycle, that is, every vertex is connected to its two first neighbors and neighbor's neighbors [1], and exact two-point resistance of the complete graph minus N edges that connect the opposite vertices [2]. In this paper, we first test these techniques by obtaining the Green's function of the one-dimensional graph with free boundaries, this is the building block for the two-point resistance of this graph, and the two-point resistance of the N -cycle graph [3]. The same techniques enabled us to obtain a closed formula for certain trigonometrical sum that appear in the work of McCoy and Orrick on the chiral Potts model [4], which was derived by Gervois and Mehta [5], using certain recursion relations in which the trigonometrical sum connected with the chiral Potts model is a special case. We have also considered the general case studied by Gervois and Mehta [5], and by Berndt and Yeap [6], here, our results agree with those in [5]. It is interesting to note that the same expression for the two-point resistance of the N -cycle graph appeared also in connection with a particle executing simple random walk on the the N -cycle [7], and the chiral Potts model [4]. having checked that our method works, then, we used it to compute other class of trigonometrical sums, some of which are related to number theory. It is interesting to point out that in all the computations of the trigonometrical sums in this paper are based on a formula by Schwatt [8] on trigonometrical power sums and the representation of the binomial coefficients by the residue operator. The Schwatt's formula is modified slightly, only in the case of the non-trivial trigonometrical sum given by $F_1(N, l, 2)$, and $F_1(N, l, 2)$, see section 4, this was also the case in our previous papers [1], [2]. This paper is organized as follows; in section 2, we give an explicit computations of the two-point resistance of the N -cycle graph and the Green's function of the one-dimensional lattice, and in section 3, we use similar techniques as in section 2, to obtain a closed formula for a trigonometrical sum arising in the chiral Potts model, also the its generalization studied by Gervois and Mehta [5] is considered and given a partial closed form. In section 4, we consider other class of trigonometrical sums which differ from those of the previous sections, some of which are related to number theory, and finally, in section 5, our conclusions are given.

2 The two-point resistance of one-dimensional lattice using the residue operator

First, let us start with the two point resistance of the N -cycle graph computations, then, move to the trigonometrical sum related to the two-point resistance of the one-dimensional lattice with free boundaries, the path graph. The two-point resistance of the N -cycle graph between any two nodes α and β is given by the following trigonometrical sum [3],

$$R(|\alpha - \beta|) = R(l) = \frac{1}{N} \sum_{n=1}^{N-1} \frac{\sin^2(nl\pi/N)}{\sin^2(n\pi/N)}. \quad (1)$$

By using the following trigonometrical identity

$$\cos(2ln\pi/N) = \sum_{s=0}^l (-1)^s \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \sin^{2s}(n\pi/N),$$

then, the above trigonometrical sum may be rewritten as

$$R(l) = \frac{1}{2N} \sum_{s=1}^l (-1)^{s+1} \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \sum_{n=1}^{N-1} \sin^{2(s-1)}(n\pi/N). \quad (2)$$

Now, from Schwatt's formula for trigonometrical power sums [8], one has

$$\sum_{n=1}^{N-1} \sin^{2(s-1)}(n\pi/N) = \frac{1}{2^{2(s-1)-1}} \sum_{t=1}^{s-1} (-1)^{t+1} \binom{2(s-1)}{s-1-t} + \frac{N-1}{2^{2(s-1)}} \binom{2(s-1)}{s-1}. \quad (3)$$

In order to evaluate $R(l)$, we use the residue operator which played a crucial role in evaluating combinatorial sums and proving combinatorial identities [9]. First, let us recall the definition of the residue operator res . To that end, let $G(w) = \sum_{k=0}^{\infty} a_k w^k$ be a generating function for a sequence $\{a_k\}$. Then the k -th coefficient of $G(w)$ may be represented by the formal residue as follows

$$a_k = \text{res}_w G(w) w^{-k-1}.$$

This is equivalent to the Cauchy integral representation of a_k ,

$$a_k = \frac{1}{2\pi i} \oint_{|z|=\rho} \frac{G(w)}{w^{k+1}} dw,$$

for coefficients of the Taylor series in a punctured neighborhood of zero. In particular, the generating function of the binomial coefficient sequence $\binom{n}{k}$ for a fixed n is given by

$$G(w) = \sum_{k=0}^n \binom{n}{k} w^k = (1+w)^n,$$

and hence

$$\binom{n}{k} = \text{res}_w (1+w)^n w^{-k-1}.$$

Before finishing this brief summary, we should mention one important property of the residue operator res , namely linearity, this is a crucial in doing computations, linearity states that given some constants α and β , then

$$\alpha \text{res}_w G_1(w) w^{-k-1} + \beta \text{res}_w G_2(w) w^{-k-1} = \text{res}_w (\alpha G_1(w) + \beta G_2(w)) w^{-k-1}.$$

Let us now, evaluate the first term in Eq.(2) using the residue operator, namely the following term

$$\begin{aligned}
R_1(l) : &= \frac{2}{N} \sum_{s=1}^l (-1)^s \frac{2l}{l+s} \binom{l+s}{l-s} \sum_{t=1}^{s-1} (-1)^t \binom{2(s-1)}{s-1-t} \\
&= \frac{2}{N} \sum_{s=1}^l (-1)^s \frac{2l}{l+s} \binom{l+s}{l-s} \sum_{t=1}^{s-1} (-1)^t \operatorname{res} \frac{(1+w)^{2(s-1)}}{w^{s-t}} \\
&= \frac{2}{N} \sum_{s=1}^l (-1)^s \frac{2l}{l+s} \binom{l+s}{l-s} \operatorname{res}_{w=0} \frac{(1+w)^{2(s-1)}}{(1+w)^3 w^{s-1}}.
\end{aligned} \tag{4}$$

In obtaining Eq.(4), we discarded an analytic term at the pole $w = 0$ of order $s - 1$. By making a change of variable $l - s = k$, then, Eq. (4), may be rewritten as

$$\begin{aligned}
R_1(l) &= (-1)^{l+1} \frac{2}{N} \operatorname{res}_{w=0} \frac{w}{(1+w)^3} \sum_{k=1}^{l-1} (-1)^k \frac{2l}{2l-k} \binom{2l-k}{k} \left(\frac{1+w}{\sqrt{w}} \right)^{2(l-k)} \\
&= (-1)^{l+1} \frac{2}{N} \operatorname{res}_{w=0} \frac{w}{(1+w)^3} \left(C_{2l} \left(\frac{1+w}{\sqrt{w}} \right) - (-1)^l \right),
\end{aligned} \tag{5}$$

where

$$C_{2l}(x) = 2T_{2l}(x/2) = \sum_{k=0}^{l-1} (-1)^k \frac{2l}{2l-k} \binom{2l-k}{k} x^{2l-2k},$$

is the normalized Chebyshev polynomial of the first kind [10], and

$$T_{2l}(x/2) = \frac{1}{2} \left[\left(\frac{x}{2} + \sqrt{(x/2)^2 - 1} \right)^{2l} + \left(\frac{x}{2} - \sqrt{(x/2)^2 - 1} \right)^{2l} \right].$$

Using the fact that $C_{2l}(\frac{1+w}{\sqrt{w}}) = \frac{1}{w^l} + w^l$, then, the final expression for the first term $R_1(l)$, reads

$$\begin{aligned}
R_1(l) &= (-1)^{l+1} \frac{2}{N} \operatorname{res}_{w=0} \frac{1}{(1+w)^3 w^{l-1}} \\
&= -\frac{l(l-1)}{N}
\end{aligned} \tag{6}$$

Similarly, the second term may written as

$$\begin{aligned}
R_2(l) : &= \frac{(N-1)}{N} \sum_{s=1}^l (-1)^s \frac{2l}{l+s} \binom{l+s}{l-s} \binom{2(s-1)}{s-1} \\
&= (-1)^{l+1} \frac{(N-1)}{N} \operatorname{res}_{w=0} \frac{1}{(1+w)^2 w^l} \\
&= \frac{l(N-1)}{N}.
\end{aligned} \tag{7}$$

Adding the contributions given by Eqs. (6), and (7), then, we get the well known formula

$$R(l) = \frac{1}{N} \sum_{n=1}^{N-1} \frac{\sin^2(nl\pi/N)}{\sin^2(n\pi/N)} = \frac{l(N-l)}{N}. \quad (8)$$

Now, we want to evaluate the following trigonometrical sum

$$F_N(l) = \frac{1}{N} \sum_{n=1}^{N-1} \frac{1 - \cos nl\pi/N}{1 - \cos n\pi/N},$$

this sum arises in connection with the two-point resistance of a path graph [3]. The evaluation of this term may be done as follows;

$$\begin{aligned} F_N(l) &= \frac{1}{N} \sum_{n=1}^{N-1} \frac{1 - \cos nl\pi/N}{1 - \cos n\pi/N} \\ &= \frac{1}{N} \sum_{n=1}^{\frac{N}{2}} \frac{1 - \cos(2n-1)l\pi/N}{1 - \cos(2n-1)\pi/N} + \frac{1}{N} \sum_{n=1}^{\frac{N}{2}-1} \frac{1 - \cos 2nl\pi/N}{1 - \cos 2n\pi/N}, \end{aligned} \quad (9)$$

where, we have assumed that N is even, similar steps may be used for N odd. It is interesting to note that in evaluating $F_N(l)$, we only need to compute the first term since the second term is related to the two-point resistance of the N -cycle graph given by Eq. (8). Then, the first term may be written as

$$\begin{aligned} \frac{1}{N} \sum_{n=1}^{\frac{N}{2}} \frac{1 - \cos(2n-1)l\pi/N}{1 - \cos(2n-1)\pi/N} &= \frac{1}{N} \sum_{n=1}^{\frac{N}{2}} \frac{\sin^2(2n-1)l\pi/2N}{\sin^2(2n-1)\pi/2N} \\ &= \frac{1}{2N} \sum_{s=1}^l (-1)^{s+1} \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \sum_{n=1}^{\frac{N}{2}} \sin^{2(s-1)} \frac{(2n-1)\pi}{2N}. \end{aligned} \quad (10)$$

By using the identity

$$\sum_{n=1}^{\frac{N}{2}} \sin^{2(s-1)}(2n-1)\pi/2N = \frac{1}{2} \left(\sum_{n=1}^{N-1} \sin^{2(s-1)} n\pi/2N + \sum_{n=1}^{N-1} (-1)^{n-1} \sin^{2(s-1)} n\pi/2N \right),$$

and the formulas for trigonometrical power sums given in [8], then, one can show

$$\sum_{n=1}^{\frac{N}{2}} \sin^{2(s-1)}(2n-1)\pi/2N = \frac{2N}{2^{2s}} \binom{2(s-1)}{s-1}, \quad (11)$$

which in turn, implies that the formula for the first term should be

$$\begin{aligned} \frac{1}{N} \sum_{n=1}^{\frac{N}{2}} \frac{1 - \cos(2n-1)l\pi/N}{1 - \cos(2n-1)\pi/N} &= \frac{1}{2} \sum_{s=1}^l (-1)^{s+1} \frac{2l}{l+s} \binom{l+s}{l-s} \binom{2(s-1)}{s-1} \\ &= \frac{l}{2}. \end{aligned} \quad (12)$$

To compute the second term given in Eq. (9), we use the following symmetry enjoyed by the two-point resistance of the N -cycle graph, N even

$$\frac{1}{N} \sum_{n=1}^{N-1} \frac{1 - \cos 2nl\pi/N}{1 - \cos 2n\pi/N} = \frac{2}{N} \sum_{n=1}^{\frac{N}{2}-1} \frac{1 - \cos 2nl\pi/N}{1 - \cos 2n\pi/N} + \frac{1}{2N} (1 - (-1)^l). \quad (13)$$

Therefore, the second term may be obtained to give the following closed formula for $F_N(l)$

$$F_N(l) = \frac{1}{N} \sum_{n=1}^{N-1} \frac{1 - \cos nl\pi/N}{1 - \cos n\pi/N} = l - \frac{1}{N} \left(\frac{l^2}{2} + \frac{1}{4} (1 - (-1)^l) \right) \quad (14)$$

3 A trigonometrical sum arising in the chiral Potts model

Here, in this section the trigonometrical sum $T_4(l) := \sum_{n=1}^{N-1} \frac{\sin^2(nl\pi/N)}{\sin^4(n\pi/N)}$ is evaluated in a closed form using the residue operator. We also give an almost closed formula for the general case $T_{2m}(l) := \sum_{n=1}^{N-1} \frac{\sin^2(nl\pi/N)}{\sin^{2m}(n\pi/N)}$, for $m \geq 1$. The first trigonometrical sum arises in the work of McCoy and Orrick on the chiral Potts model [4], this sum with other trigonometrical identities were proved by Gervois and Mehta [5]. the second sum was considered by Gervois and Mehta [5] using a recursion formula satisfied by $T_{2m}(l)$. It was shown by Berndt and Yeap [6] that the second sum may be written in terms of sums of the Bernoulli numbers. By following the same techniques as in the previous section, the above first sum may be written as

$$\begin{aligned} T_4(l) &= \sum_{n=1}^{N-1} \frac{\sin^2(nl\pi/N)}{\sin^4(n\pi/N)} \\ &= l^2 \sum_{n=1}^{N-1} \frac{1}{\sin^2(n\pi/N)} + \frac{1}{2} \sum_{s=2}^l (-1)^{s+1} \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \sum_{n=1}^{N-1} \sin^{2(s-2)}(n\pi/N) \\ &= \frac{l^2}{3} (N^2 - 1) + 8 \sum_{s=2}^l (-1)^s \frac{2l}{l+s} \binom{l+s}{l-s} \sum_{t=1}^{s-2} (-1)^t \binom{2(s-2)}{s-2-t} \\ &\quad + 4(N-1) \sum_{s=2}^l (-1)^{s+1} \frac{2l}{l+s} \binom{l+s}{l-s} \binom{2(s-2)}{s-2}, \end{aligned} \quad (15)$$

the first term in the above equation follows from the well known identity

$$\sum_{n=1}^{N-1} \frac{1}{\sin^2(n\pi/N)} = \frac{N^2 - 1}{3},$$

while the second and the third terms may be computed using the residue operator as in the previous section to give,

$$\begin{aligned} \sum_{s=2}^l (-1)^{s+1} \frac{2l}{l+s} \binom{l+s}{l-s} \sum_{t=1}^{s-2} (-1)^t \binom{2(s-2)}{s-2-t} &= (-1)^{l+1} \text{res}_{w=0} \frac{1}{(1+w)^5 w^{l-2}} \\ &= \frac{1}{4!} (l+1)l(l-1)(l-2), \end{aligned} \quad (16)$$

and

$$\begin{aligned} \sum_{s=2}^l (-1)^{s+1} \frac{2l}{l+s} \binom{l+s}{l-s} \binom{2(s-2)}{s-2} &= (-1)^{l+1} \text{res}_{w=0} \frac{1}{(1+w)^4 w^{l-1}} \\ &= -\frac{1}{3!} (l+1)l(l-1). \end{aligned} \quad (17)$$

Therefore, the closed formula for the sum given in Eq. (15), reads

$$\begin{aligned} \sum_{n=1}^{N-1} \frac{\sin^2(nl\pi/N)}{\sin^4(n\pi/N)} &= \frac{l^2}{3} (N^2 - 1) + \frac{1}{3} (l+1)l(l-1)(l-2) - \frac{2(N-1)}{3} (l+1)l(l-1) \\ &= \frac{l^2}{3} (N-l)(N+l) + \frac{2}{3} (N-l). \end{aligned} \quad (18)$$

This is exactly the result obtained by Gervois and Mehta using a recursion formula satisfied by $T_{2m}(l)$ [5]. Next, we will give another formula for the sum $T_{2m}(l)$ that differs from those given in [5] and [6], and contains two terms one of which has a closed form. Now, $T_{2m}(l)$, may be written as

$$\begin{aligned} T_{2m}(l) &= \sum_{n=1}^{N-1} \frac{\sin^2(nl\pi/N)}{\sin^{2m}(n\pi/N)} \\ &= \frac{1}{2} \sum_{n=1}^{N-1} \sum_{s=1}^{m-1} (-1)^{s+1} \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \frac{1}{\sin^{2(m-s)}(n\pi/N)} \\ &\quad + \frac{1}{2} \sum_{s=m}^l (-1)^{s+1} \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \sum_{n=1}^{N-1} \sin^{2(s-m)}(n\pi/N). \end{aligned} \quad (19)$$

The first term on the right hand side $T_{2k} =: \sum_{n=1}^{N-1} \frac{1}{\sin^{2k}(n\pi/N)}$, may be computed using a recursion formula given in [5], while the second term as it will be shown shortly, has a closed

form formula. The second may be written as follows

$$\begin{aligned}
\tilde{T}_{2m}(l) : &= \frac{1}{2} \sum_{s=m}^l (-1)^{s+1} \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \sum_{n=1}^{N-1} \sin^{2(s-m)}(n\pi/N) \\
&= 2^{2m-1} \sum_{s=m}^l (-1)^{s+1} \frac{2l}{l+s} \binom{l+s}{l-s} \sum_{t=1}^{s-m} (-1)^t \binom{2(s-m)}{s-m-t} \\
&+ 2^{2m-2} (N-1) \sum_{s=m}^l (-1)^{s+1} \frac{2l}{l+s} \binom{l+s}{l-s} \binom{2(s-m)}{s-m} \\
&= (-1)^{l+1} 2^{2m-1} \text{res}_{w=0} \frac{1}{(1+w)^{2m+1} w^{l-m}} \\
&+ (-1)^{l+1} 2^{2m-2} (N-1) \text{res}_{w=0} \frac{1}{(1+w)^{2m} w^{l+1-m}} \\
&= (-1)^m 2^{2m-1} \frac{(l+m-1)!}{(l-m-1)!(2m)!} + (-1)^{m+1} (N-1) 2^{2m-2} \frac{(l+m-1)!}{(l-m)!(2m-1)!}
\end{aligned} \tag{20}$$

Therefore, we succeeded in writing $\tilde{T}_{2m}(l)$ in a closed form formula, One can check easily that our results agree with those given in [5], and so the formula for $T_{2m}(l)$ becomes

$$\begin{aligned}
T_{2m}(l) &= \sum_{n=1}^{N-1} \frac{\sin^2(nl\pi/N)}{\sin^{2m}(n\pi/N)} \\
&= \frac{1}{2} \sum_{n=1}^{N-1} \sum_{s=1}^{m-1} (-1)^{s+1} \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \frac{1}{\sin^{2(m-s)}(n\pi/N)} \\
&+ (-1)^{m+1} 2^{2m-1} \frac{(l+m-1)!}{(l-m)!(2m)!} (mN-l).
\end{aligned} \tag{21}$$

setting $m = 1, 2$, then, our previous results given by Eq's. (8) and (18) respectively are recovered. From Eq. (21), it is clear that in order to have a closed formula for $T_{2m}(l)$, one needs also, the exact expressions for T_{2k} , $k = 1 \cdots, m-1$. The expression for T_{2m} may be obtained from $T_{2m}(l)$ from the following simple formula;

$$\begin{aligned}
\sum_{l=1}^{N-1} T_{2m}(l) &= \sum_{l=1}^{N-1} \sum_{n=1}^{N-1} \frac{\sin^2(nl\pi/N)}{\sin^{2m}(n\pi/N)} \\
&= \frac{N}{2} \sum_{n=1}^{N-1} \frac{1}{\sin^{2m}(n\pi/N)},
\end{aligned} \tag{22}$$

Therefore, we may write

$$\begin{aligned}
T_{2m} &= \sum_{l=1}^{N-1} \frac{1}{\sin^{2m}(n\pi/N)} \\
&= \frac{1}{N} \sum_{l=1}^{N-1} \sum_{n=1}^{N-1} \sum_{s=1}^{m-1} (-1)^{s+1} \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \frac{1}{\sin^{2(m-s)}(n\pi/N)} \\
&+ \frac{2}{N} \sum_{l=1}^{N-1} (-1)^{l+1} 2^{2m-1} \operatorname{res}_{w=0} \frac{1}{(1+w)^{2m+1} w^{l-m}} \\
&+ \frac{2}{N} \sum_{l=1}^{N-1} (-1)^{l+1} 2^{2m-2} (N-1) \operatorname{res}_{w=0} \frac{1}{(1+w)^{2m} w^{l+1-m}} \\
&= \frac{1}{N} \sum_{l=1}^{N-1} \sum_{n=1}^{N-1} \sum_{s=1}^{m-1} (-1)^{s+1} \frac{l}{l+s} \binom{l+s}{l-s} 2^{2s} \frac{1}{\sin^{2(m-s)}(n\pi/N)} \\
&+ (-1)^{m+1} 2^{2m-1} \frac{(N+m-1)!}{N(N-m-1)!(2m+1)!} (2mN+1-N). \tag{23}
\end{aligned}$$

This is another recursion formula that relates the different T_{2m} 's, for $m=1$ the first term in Eq. (23), does not contribute and the second term gives the well known formula $T_2 = \frac{N^2-1}{3}$. Now, for $m=2$, the first term may be computed to give $\frac{(N^2-1)(N-1)(2N-1)}{9}$, while the second term gives $-\frac{(N^2-1)(N-2)(3N+1)}{15}$, and hence, $T_4 = \frac{(N^2-1)(N^2+11)}{45}$ in a full agreement with [5], [6].

4 Some trigonometrical sums related to number theory

In this section other trigonometrical sums will be considered using similar techniques as in the previous sections. we will start with the following sum

$$S(l) := \sum_{n=1}^{N-1} (-1)^n \frac{\sin^2(nl\pi/N)}{\sin^2(n\pi/N)},$$

which is the alternating sum associated with the sum $R(l)$ given in Eq. (1). To obtain a closed formula for this sum, we follow basically the same computation carried out for $R(l)$, except this time, the sum over n is non-vanishing only for N even

$$\sum_{n=1}^{N-1} (-1)^n \sin^{2(s-1)}(n\pi/N) = \frac{1}{2^{2(s-1)-1}} \sum_{t=1}^{s-1} (-1)^{t+1} \binom{2(s-1)}{s-1-t} + \frac{-1}{2^{2(s-1)}} \binom{2(s-1)}{s-1}, \tag{24}$$

Comparing Eq. (24) and Eq. (3), and using the previous results, then, without any further computations one has the following simple formula

$$S(l) = \sum_{n=1}^{N-1} (-1)^n \frac{\sin^2(nl\pi/N)}{\sin^2(n\pi/N)} = -l^2. \tag{25}$$

Due to the symmetry enjoyed by $S(l)$, $S(l) = S(N - l)$, the right hand side should be read with this constraint, that is for both l , and $N - l$, $S(l) = -l^2$. Next, let us consider the sums $S_1(l) := \sum_{n=1}^{N-1} \frac{\sin(nl\pi/N)}{\sin(n\pi/N)}$ and $S_2(l) := \sum_{n=1}^{N-1} (-1)^n \frac{\sin(nl\pi/N)}{\sin(n\pi/N)}$ that are closely related. By using the following formula [11],

$$\frac{\sin(nl\pi/N)}{\sin(n\pi/N)} = \sum_{s \geq 0} (-1)^s \binom{l-s-1}{s} 2^{l-2s-1} \cos^{l-2s-1}(n\pi/N), \quad (26)$$

then, for l odd, one has

$$S_1(l) = \sum_{n=1}^{N-1} \frac{\sin(2l-1)n\pi/N}{\sin n\pi/N} = \sum_{s \geq 0} (-1)^s \binom{2l-2-s}{s} 2^{2l-2-2s} \sum_{n=1}^{N-1} \cos^{2l-2-2s}(n\pi/N) \quad (27)$$

The sum over n , may be computed from Schwatt's book [8], see Eq. (107), page 221 to give

$$\sum_{n=1}^{N-1} \cos^{2l-2-2s}(n\pi/N) = \frac{-2}{2^{2l-2-2s}} \sum_{t=1}^{l-1-s} \binom{2l-2-2s}{l-1-s-t} + \frac{N-1}{2^{2l-2-2s}} \binom{2l-2-2s}{l-1-s}, \quad (28)$$

then, the first contribution to the sum given in Eq. (27) is

$$\begin{aligned} S_1(l)' &= -2 \sum_{s=0}^{l-1} (-1)^s \binom{2l-2-s}{s} \sum_{t=1}^{l-1-s} \binom{2l-2-2s}{l-1-s-t} = \\ &= -2 \operatorname{res}_{w=0} \sum_{s=0}^{l-1} (-1)^s \binom{2l-2-s}{s} \left(\frac{1+w}{\sqrt{w}} \right)^{2l-2-2s} \frac{1}{1-w} \\ &= -2 \operatorname{res}_{w=0} U_{2l-2} \left(\frac{1+w}{2\sqrt{w}} \right) \frac{1}{1-w}, \end{aligned} \quad (29)$$

in obtaining the last line of the above equation we used the expresion for the normalized Chebysheve polynomial of the second kind $U_n(\frac{x}{2}) = \sum_{k=0}^{[n/2]} (-1)^k \binom{n-k}{k} x^{n-2k}$. The residue may be evaluated using

$$U_n\left(\frac{x}{2}\right) = \frac{(x + \sqrt{x^2 - 1})^{n+1} - (x - \sqrt{x^2 - 1})^{n+1}}{2\sqrt{x^2 - 1}},$$

to obtain

$$S_1(l)' = -2 \operatorname{res}_{w=0} \frac{1}{w^{l-1}} \frac{1}{(1-w)^2} = -2(l-1). \quad (30)$$

Similarly, the second contribution reads

$$\begin{aligned} S_1(l)'' &= (N-1) \operatorname{res}_{w=0} U_{2l-2} \left(\frac{1+w}{2\sqrt{w}} \right) \frac{1}{w} \\ &= N-1. \end{aligned} \quad (31)$$

Therefore, combining these contributions, the closed formula for $S_1(l)$ is

$$S_1(l) = \sum_{n=1}^{N-1} \frac{\sin(2l-1)n\pi/N}{\sin n\pi/N} = N - (2l-1). \quad (32)$$

It is not difficult to show that there is no contribution to the sum $S_1(l)$ for l even. To carry out the computation for

$$S_2(l) = \sum_{s \geq 0} (-1)^s \binom{l-s-1}{s} 2^{l-2s-1} \sum_{n=1}^{N-1} (-1)^n \cos^{l-2s-1}(n\pi/N),$$

one has to evaluate the sum over n , this sum turns out to depend on both l, N unlike the previous case. Using formulas given by Eq (113), and Eq (114) in [8], we have

$$\sum_{n=1}^{N-1} (-1)^n \cos^{l-2s-1}(n\pi/N) = \frac{-2}{2^{2l-2-2s}} \sum_{t=1}^{l-1-s} \binom{2l-2-2s}{l-1-s-t} - \frac{1}{2^{2l-2-2s}} \binom{2l-2-2s}{s-l-1}, \quad (33)$$

for l odd, N even, and the sum vanishes for l odd, N odd. If l is even, then, the above sum is non-vanishing only for N odd

$$\sum_{n=1}^{N-1} (-1)^n \cos^{l-2s-1}(n\pi/N) = \frac{-2}{2^{2l-1-2s}} \sum_{t=0}^{l-1-s} \binom{2l-1-2s}{l-1-s-t}. \quad (34)$$

Therefore, for l odd it follows from Eq. (33) that we have

$$\begin{aligned} S_2(l) &= \sum_{n=1}^{N-1} (-1)^n \frac{\sin(nl\pi/N)}{\sin(n\pi/N)} \\ &= -2 \operatorname{res}_{w=0} \frac{1}{w^{l-1}} \frac{1}{(1-w)^2} - \operatorname{res}_{w=0} \frac{1}{w^l} \frac{1}{(1-w)} = -(2l-1), \end{aligned} \quad (35)$$

similarly, for l even, Eq. (34) gives

$$S_2(l) = \sum_{n=1}^{N-1} (-1)^n \frac{\sin(nl\pi/N)}{\sin(n\pi/N)} = -2 \operatorname{res}_{w=0} \frac{1}{w^l} \frac{1}{(1-w)^2} = -2l, \quad (36)$$

that is, $S_2(l) = -l$, for both odd and even l , these results were recently verified by simulations without proof in connection with number theory¹. It is interesting to note that the closed formula for $S_2(l)$ may be expected simply, let l go to $N-l$ in $S_1(l)$, to obtain formally $S_2(l) = -l$. Now, we will consider the following non-trivial and interesting trigonometrical sums,

$$F_1(N, l, 2) := \sum_{n=1}^{N-1} \frac{\sin(nl\pi/N)}{\sin(n\pi/N)} \frac{\sin(2nl\pi/N)}{\sin(2n\pi/N)},$$

¹Anonymous author working on characters of a finite field and the Polya-Vinogradov inequality.

and

$$F_2(N, l, 2) := \sum_{n=1}^{N-1} (-1)^n \frac{\sin(nl\pi/N)}{\sin(n\pi/N)} \frac{\sin(2nl\pi/N)}{\sin(2n\pi/N)},$$

where $F_1(N, N-l, 2) = F_2(N, l, 2)$, and N odd. So, if $F_2(N, l, 2)$ is known, then, $F_2(N, l, 2)$ may be obtained and vice-versa. It is not difficult to show that $F_1(N, 2l, 2) = 0$, and hence, we will consider only the sum $F_1(N, 2l-1, 2)$. The latter may be written as

$$\begin{aligned} F_1(N, 2l-1, 2) &= \sum_{n=1}^{N-1} \frac{\sin(n(2l-1)\pi/N)}{\sin(n\pi/N)} \frac{\sin(2n(2l-1)\pi/N)}{\sin(2n\pi/N)} \\ &= \sum_{s,k \geq 0}^{l-1} (-1)^{s+k} \binom{2l-2-s}{s} \binom{2l-2-k}{k} 2^{2(2l-2)-2(s+k)} \\ &\quad \times \sum_{j=0}^{2l-2-2k} (-1)^j 2^j \binom{2l-2-2k}{k} \sum_{n=1}^{N-1} \cos^{2l-2-2(s-j)}(n\pi/N). \end{aligned} \quad (37)$$

The sum over n , formally looks like that given in Eq. (28), however, in this case the variable t , may be a multiple of N , and in that case the Schwatt's formula given by Eq (107), does not work it has to be modified slightly. The formula that takes into account this fact may be shown to be given by

$$\begin{aligned} \sum_{n=1}^{N-1} \cos^{2l-2-2(s-j)}(n\pi/N) &= \frac{-2}{2^{2l-2-2(s-j)}} \sum_{t=1}^{l-1-s} \binom{2l-2-2(s-j)}{l-1-(s-j)-t} \\ &\quad + \frac{N-1}{2^{2l-2-2(s-j)}} \binom{2l-2-2(s-j)}{l-1-(s-j)} \\ &\quad + \frac{2N}{2^{2l-2-2(s-j)}} \sum_{p=1}^{[l-1-(s-j)/N]} \binom{2l-2-2(s-j)}{l-1-(s-j)-pN}, \end{aligned} \quad (38)$$

where the first two terms in the above formula are those expected from Eq. (28), while the third term is precisely the correction to the formula Eq. (28), in the case t is congruent to N . Therefore, there are three contributions to the sum given in Eq. (37), the first of which

reads

$$\begin{aligned}
F_1'(N, 2l-1, 2) &= -2 \sum_{s,k \geq 0}^{l-1} (-1)^{s+k} \binom{2l-2-s}{s} \binom{2l-2-k}{k} 2^{2l-2-2k} \\
&\times \sum_{j=0}^{2l-2-2k} (-1)^j \frac{1}{2^j} \binom{2l-2-2k}{k} \sum_{t=1}^{l-1-s} \binom{2l-2-2(s-j)}{l-1-(s-j)-t} \\
&= -2 \operatorname{res}_{w=0} U_{2l-2} \left(\frac{1+w}{2\sqrt{w}} \right) U_{2l-2} \left(\frac{1+w^2}{2w} \right) \frac{1}{1-w} \\
&= -2 \operatorname{res}_{w=0} \left(\frac{1}{w^{3l-3}} - \frac{1}{w^{l-2}} \right) \frac{1}{(1-w)^2} \frac{1}{1-w^2} \\
&= -\frac{1}{2} (3l-2)(3l-3) + \frac{1}{2} (l-1)(l-2) - l + \frac{1}{2} (1 - (-1)^l) \quad (39)
\end{aligned}$$

while the second contribution reads

$$\begin{aligned}
F_1''(N, 2l-1, 2) &= (N-1) \sum_{s,k \geq 0}^{l-1} (-1)^{s+k} \binom{2l-2-s}{s} \binom{2l-2-k}{k} 2^{2l-2-2k} \\
&\times \sum_{j=0}^{2l-2-2k} (-1)^j \frac{1}{2^j} \binom{2l-2-2k}{k} \binom{2l-2-2(s-j)}{l-1-(s-j)} \\
&= (N-1) \operatorname{res}_{w=0} U_{2l-2} \left(\frac{1+w}{2\sqrt{w}} \right) U_{2l-2} \left(\frac{1+w^2}{2w} \right) \frac{1}{w} \\
&= (N-1) \operatorname{res}_{w=0} \left(\frac{1}{w^{3l-2}} - \frac{1}{w^{l-1}} \right) \frac{1}{1-w^2} \frac{1}{1-w} \\
&= \frac{N-1}{2} (2l-1 - (-1)^l). \quad (40)
\end{aligned}$$

To obtain the last contribution we write the sum over p , in Eq. (38), as

$$\sum_{p=1}^{[l-1-(s-j)/N]} \binom{2l-2-2(s-j)}{l-1-(s-j)-pN} = \operatorname{res}_{w=0} \left(\frac{(1+w)^{2l-2-2(s-j)} w^N}{w^{l-(s-j)} (1-w^N)} \right),$$

and using the fact that $l \leq N - 1$, then, the third contribution may be computed to give

$$\begin{aligned}
F_1'''(N, 2l - 1, 2) &= 2N \sum_{s, k \geq 0}^{l-1} (-1)^{s+k} \binom{2l-2-s}{s} \binom{2l-2-k}{k} 2^{2l-2-2k} \\
&\times \sum_{j=0}^{2l-2-2k} (-1)^j \frac{1}{2^j} \binom{2l-2-2k}{k} \text{res}_{w=0} \left(\frac{(1+w)^{2l-2-2(s-j)} w^N}{w^{l-(s-j)} (1-w^N)} \right) \\
&= 2N \text{res}_{w=0} \left(\frac{1}{1-w^N} U_{2l-2} \left(\frac{1+w}{2\sqrt{w}} \right) U_{2l-2} \left(\frac{1+w^2}{2w} \right) w^{N-1} \right) \\
&= 2N \text{res}_{w=0} \frac{1}{1-w^N} \frac{1}{w^{3l-2-N}} \frac{1}{(1-w)(1-w^2)} \\
&= N \left(3l - 2 - N + \frac{1}{2} (1 - (-1)^{l-N}) \right). \tag{41}
\end{aligned}$$

Note that this will contribute only for $3l - 2 > N$, and as a consequence the closed formula for the sum $F_1(N, 2l - 1, 2)$ is

$$\begin{aligned}
F_1(N, 2l - 1, 2) &= \sum_{n=1}^{N-1} \frac{\sin(n(2l-1)\pi/N)}{\sin(n\pi/N)} \frac{\sin(2n(2l-1)\pi/N)}{\sin(2n\pi/N)} \\
&= -\frac{1}{2} (3l-2)(3l-3) + \frac{1}{2} (l-1)(l-2) - l + \frac{1}{2} (1 - (-1)^l) \\
&+ \frac{N-1}{2} \left(2l-1 - (-1)^l \right) + N \left(3l-2-N + \frac{1}{2} (1 - (-1)^{l-N}) \right). \tag{42}
\end{aligned}$$

Having obtained a closed formula for the sum $F_1(N, 2l-1, 2)$, we turn now to the alternating sum

$$F_2(N, l, 2) = \sum_{n=1}^{N-1} (-1)^n \frac{\sin(nl\pi/N)}{\sin(n\pi/N)} \frac{\sin(2nl\pi/N)}{\sin(2n\pi/N)},$$

in this case, for N odd, it is easy to show that the sum is non-vanishing only for l is even. The computation is basically the same, the only difference is that in this case we have an alternating sum and l is even

$$\begin{aligned}
F_2(N, 2l, 2) &= \sum_{n=1}^{N-1} (-1)^n \frac{\sin((2l)n\pi/N)}{\sin(n\pi/N)} \frac{\sin((2l)2n\pi/N)}{\sin(2n\pi/N)} \\
&= \sum_{s, k \geq 0}^{[(2l-1)/2]} (-1)^{s+k} \binom{2l-1-s}{s} \binom{2l-1-k}{k} 2^{2(2l-1)-2(s+k)} \\
&\times \sum_{j=0}^{2l-1-2k} (-1)^j 2^j \binom{2l-1-2k}{k} \sum_{n=1}^{N-1} (-1)^n \cos^{2l-1-2(s-j)}(n\pi/N). \tag{43}
\end{aligned}$$

The sum over n may be carried out using Eq. (114), in [8] with the slight modification as explained before, then, it is not difficult to show

$$\begin{aligned} \sum_{n=1}^{N-1} (-1)^n \cos^{2l-1-2(s-j)-1}(n\pi/N) &= \frac{-2}{2^{2l-1-2(s-j)}} \sum_{t=0}^{l-1-s} \binom{2l-1-2(s-j)}{l-1-(s-j)-t} \\ &+ \frac{2N}{2^{2l-1-2(s-j)}} \sum_{p \geq 1} \binom{2l-1-2(s-j)}{l-1-(s-j)-\frac{(2p-1)N-1}{2}}, \end{aligned} \quad (44)$$

where the sum over p , may be written as

$$\sum_{p \geq 1} \binom{2l-1-2(s-j)}{l-1-(s-j)-\frac{(2p-1)N-1}{2}} = \text{res}_{w=0} \left(\frac{(1+w)^{2l-1-2(s-j)} w^{N/2}}{w^{l-(s-j)+1/2} (1-w^N)} \right).$$

By using Eq. (43), computations show that the closed formula for the sum $F_2(N, 2l, 2)$ is

$$\begin{aligned} F_2(N, 2l, 2) &= \sum_{n=1}^{N-1} (-1)^n \frac{\sin((2l)n\pi/N)}{\sin(n\pi/N)} \frac{\sin((2l)2n\pi/N)}{\sin(2n\pi/N)} \\ &= -2 \text{res}_{w=0} \left(U_{2l-2} \left(\frac{1+w}{2\sqrt{w}} \right) U_{2l-2} \left(\frac{1+w^2}{2w} \right) \frac{1}{\sqrt{w}(1-w)} \right) \\ &+ 2N \text{res}_{w=0} \frac{1}{1-w^N} \left(U_{2l-1} \left(\frac{1+w}{2\sqrt{w}} \right) U_{2l-1} \left(\frac{1+w^2}{2w} \right) \frac{w^{N/2}}{w} \right) \\ &= -2 \text{res}_{w=0} \left(\frac{1}{w^{3l-1}} - \frac{1}{w^{l-1}} \right) \frac{1}{(1-w)^2} \frac{1}{1-w^2} \\ &+ 2N \text{res}_{w=0} \frac{1}{1-w^N} \frac{1}{w^{3l-(N+1)/2}} \frac{1}{(1-w)(1-w^2)} \\ &= -\frac{1}{2} 3l(3l-1) + \frac{1}{2} l(l-1) - l + N \left(3l - \frac{(N+1)}{2} + \frac{1}{2} (1 - (-1)^{l-\frac{(N+1)}{2}}) \right), \end{aligned} \quad (45)$$

where the last term whose coefficient is N , contributes only for $3l > \frac{(N+1)}{2}$. Let us now, check that the formulas $F_1(N, 2l-1, 2)$, $F_2(N, 2l, 2)$ are consistent with symmetry discussed earlier, that is, $F_1(N, N-l, 2) = F_2(N, l, 2)$, this in turns implies that the correctness of the formulas. To do so, we will give some explicit examples, from the expression of $F_1(N, 2l-1, 2)$ given in Eq. (42), it is clear that the sum should be $N-1$, for $l=1$ and to check this, one has to take into account that when substituting $l=1$ in the formula, the last term of Eq. (42) does not contribute. From the symmetry that relates the two sums, we should have $F_2(N, N-1, 2) = N-1$. Indeed, this is the case, we simply let $l = \frac{N-1}{2}$ into Eq. (45), this time, however, the last term of this equation does contribute. An explicit computation shows that $F_2(N, 2, 2) = -4$, for $N > 3$, and $F_2(N, 2, 2) = 2$, for $N = 3$, it is interesting to note that these two cases for $l=1$ are contained in the last term of Eq. (44),

since for $N > 3$, the last term is equal to 0, and hence $F_2(N, 2, 2) = -4$, while for $N = 3$, the last term is equal to 6, that is, our formula gives the right answer. Using the symmetry, we obtain $F_1(N, N - 2, 2) = -4$, this can be easily checked using our formula given by Eq. (42), and $l = \frac{N-1}{2}$.

5 Conclusion

To conclude, in this paper we were able to obtain a number of closed formulas for trigonometrical sums that appear in one-dimensional lattice, the chiral Potts model and number theory. Here, we followed similar formalism developed for the circulant electrical network (circulant graphs) [1], [2], as a consequence the non-trivial circulant electrical networks (the cycle and complete graphs are not included) are related to non-trivial trigonometrical sums in number theory. For example in [2], we had to introduce certain numbers that we called the Bejaia and Pisa numbers with well known properties so that the trigonometrical sums that arise are written in terms of these numbers nicely. By using the well known connection between the electrical networks and the random walks [12], one may hope to give interpretations to some of the trigonometrical sums in number theory other than those associated with the two-point resistance of a given electrical network, since the latter provides an alternative way to compute the basic quantity relevant to random walks known as the first passage time, the expected time to hit a target node for the first time for a walker starting from a source node [13].

References

- [1] N.Chair submitted
- [2] N.Chair submitted, arXiv:1206.1519v1 [math-ph]
- [3] F. Y. Wu, J. Phys. A: Math. Gen. 37 6653 (2004)
- [4] Barry M. McCoy, William P. Orrick, J.Statist.Phys. 83, 839 (1996)
- [5] A. Gervois and M. L. Mehta J. Math. Phys. 36, 5098 (1995)
- [6] B. C. Berndt and B. P. Yeap, Adv. in Appl. Math. 29, 358 (2002)
- [7] D.J Aldous, J.Fill, 2012. Reversible Markov chains and random walks on graphs. Available in: <http://www.stat.berkeley.edu/~aldous/RWG/book.html>
- [8] I. J. Schwatt, An Introduction to the Operations with Series. Philadelphia, press of the university of Pennsylvania, (1924).
- [9] G. P. Egorychev. Integral representation and the computation of combinatorial sums American Mathematical Soc, 1984.
- [10] T. J Rivlin, . Chebyshev Polynomials. New York: Wiley, 1990.
- [11] E. W Hobson A treatise on plane and advanced trigonometry, seventh edition Dover Publications, Inc. New York. New York
- [12] P.G. Doyle, J.L. Snell, Random Walks and Electrical Networks, The Mathematical Association of America, Washington, DC, 1984
- [13] Tetali, P. (1991). Journal of Theoretical Probability, 4, 101 (1991)